


Probability and Statistics

Nimisha Roy

Lecturer, SCI, College of Computing, Georgia Tech

Director, Online Undergraduate Initiatives

Outline

- Probability Distributions 
- Joint and Conditional Probability Distributions
- Bayes' Rule
- Mean and Variance
- Properties of Gaussian Distribution
- Maximum Likelihood Estimation

Probability

- A **sample space S** is the set of all possible outcomes of a conceptual or physical, repeatable experiment. (S can be finite or infinite.)
 - E.g., S may be the set of all possible outcomes of a dice roll: S
(1 2 3 4 5 6)
 - E.g., S may be the set of all possible nucleotides of a DNA site: S
(*A C G T*)
- E.g., S may be the set of all possible time-space positions of an aircraft on a radar screen.
- An **Event A** is any subset of S
 - Seeing "1" or "6" in a dice roll; observing a "G" at a site; UA007 in space-time interval



Three Key Ingredients in Probability Theory

A **sample space** is a collection of all possible **outcomes**

Random variables X represents **outcomes** in sample space

Probability of a random variable to happen $p(x) = p(X = x)$

$$p(x) \geq 0$$

Continuous variable

Definition: Takes values from a continuous range (e.g., any real number within an interval).

Distribution: Governed by a **probability density function** (PDF).

Key Concepts:

- The **density** represents likelihood, but not actual probability at a specific point.
- Example: **Temperature**, which can be any real value (e.g., 72.3°F).
- Common distribution: **Gaussian (Normal) Distribution**.

$$\int_x p(x) dx = 1$$

Discrete variable

Definition: Takes values from a countable set (e.g., integers).

Distribution: Governed by a **probability mass function** (PMF).

Key Concepts:

- The function directly gives **probability values**.
- Example: **A coin flip** (e.g., 0 for tails, 1 for heads).
- Common distribution: **Bernoulli Distribution**.

$$\sum_{x \in A} p(x) = 1$$

Continuous Probability Functions

- Examples:

- Uniform Density Function:

$$f_x(x) = \begin{cases} \frac{1}{b-a} & \text{for } a \leq x \leq b \\ 0 & \text{otherwise} \end{cases}$$

- Exponential Density Function:

$$f_x(x) = \frac{1}{\mu} e^{-\frac{x}{\mu}} \quad \text{for } x \geq 0$$

- Gaussian(Normal) Density Function

$$f_x(x) = \frac{1}{\sqrt{2\pi}\sigma} e^{-\frac{(x-\mu)^2}{2\sigma^2}}$$

Discrete Probability Functions

- Examples:

- Bernoulli Distribution:

- $$\begin{cases} 1 - p & \text{for } x = 0 \\ p & \text{for } x = 1 \end{cases}$$

In Bernoulli, just a **single** trial is conducted

- Binomial Distribution:

- $$P(X = k) = \binom{n}{k} p^k (1 - p)^{n-k}$$

k is number of successes

n-k is number of failures

$\binom{n}{k}$ The total number of ways of selection **k** distinct combinations of **n** trials, **irrespective of order**.

E.g. - Given a biased coin with $\mu = 0.3$ for tail ($x = 1$), what is the probability of getting 4 heads given we flip the coin 10 times?

Outline

- Probability Distributions
- Joint and Conditional Probability Distributions
- Bayes' Rule
- Mean and Variance
- Properties of Gaussian Distribution
- Maximum Likelihood Estimation



Recap

- Sample Space and Event
- Probability Distribution Function
- Objective Function
- Joint Probability, Marginal Probability & Conditional Probability

Example



X = Throw a dice



Y = Flip a coin

X and **Y** are random variables

N = total number of trials

n_{ij} = Number of occurrence

		X						C_j
		$x_{i=1} = 1$	$x_{i=2} = 2$	$x_{i=3} = 3$	$x_{i=4} = 4$	$x_{i=5} = 5$	$x_{i=6} = 6$	
Y	$y_{j=2} = tail$	$n_{ij} = 3$	$n_{ij} = 4$	$n_{ij} = 2$	$n_{ij} = 5$	$n_{ij} = 1$	$n_{ij} = 5$	20
	$y_{j=1} = head$	$n_{ij} = 2$	$n_{ij} = 2$	$n_{ij} = 4$	$n_{ij} = 2$	$n_{ij} = 4$	$n_{ij} = 1$	15
C_i		5	6	6	7	5	6	N=35

X
 $x_{i=1} = 1$ $x_{i=2} = 2$ $x_{i=3} = 3$ $x_{i=4} = 4$ $x_{i=5} = 5$ $x_{i=6} = 6$
 C_j **Y** $y_{j=2} = tail$ $y_{j=1} = head$ C_i

$n_{ij} = 3$	$n_{ij} = 4$	$n_{ij} = 2$	$n_{ij} = 5$	$n_{ij} = 1$	$n_{ij} = 5$	20
$n_{ij} = 2$	$n_{ij} = 2$	$n_{ij} = 4$	$n_{ij} = 2$	$n_{ij} = 4$	$n_{ij} = 1$	15
5	6	6	7	5	6	N=35

Probability:

$$p(X = x_i) = \frac{c_i}{N}$$

Joint probability:

$$p(X = x_i, Y = y_j) = \frac{n_{ij}}{N}$$

Conditional probability:

$$p(Y = y_j | X = x_i) = \frac{n_{ij}}{c_i}$$

Sum rule

$$p(X = x_i) = \sum_{j=1}^L p(X = x_i, Y = y_j) \Rightarrow p(X) = \sum_Y P(X, Y)$$

Product rule

$$p(X = x_i, Y = y_j) = \frac{n_{ij}}{N} = \frac{n_{ij}}{c_i} \frac{c_i}{N} = p(Y = y_j | X = x_i) p(X = x_i)$$

$$p(X, Y) = p(Y|X)p(X)$$

Conditional Independence

- Examples:

$$P(\text{Virus} | \text{DrinkBeer}) = P(\text{Virus})$$

iff **Virus** is independent of **Drink Beer**

$$P(\text{Flu} | \text{Virus}, \text{DrinkBeer}) = P(\text{Flu} | \text{Virus})$$

iff **Flu** is independent of **Drink Beer**, given **Virus**


$$\begin{aligned} P(\text{Headache} | \text{Flu}, \text{Virus}, \text{DrinkBeer}) \\ = P(\text{Headache} | \text{Flu}, \text{DrinkBeer}) \end{aligned}$$

iff **Headache** is independent of **Virus**, given **Flu** and **Drink Beer**

Assume the above independence, we obtain:

$$\begin{aligned} P(\text{Headache}, \text{Flu}, \text{Virus}, \text{DrinkBeer}) \\ = P(\text{Headache} | \text{Flu}, \text{Virus}, \text{DrinkBeer}) P(\text{Flu} | \text{Virus}, \text{DrinkBeer}) \\ P(\text{Virus} | \text{DrinkBeer}) P(\text{DrinkBeer}) \\ = P(\text{Headache} | \text{Flu}, \text{DrinkBeer}) P(\text{Flu} | \text{Virus}) P(\text{Virus}) P(\text{DrinkBeer}) \end{aligned}$$

Outline

- Probability Distributions
- Joint and Conditional Probability Distributions
- Bayes' Rule 
- Mean and Variance
- Properties of Gaussian Distribution
- Maximum Likelihood Estimation

Bayes' Rule

- $P(X|Y)$ = Fraction of the worlds in which X is true given that Y is also true.
- For example:
 - H = "Having a headache"
 - F = "Coming down with flu"
 - $P(\text{Headache}|\text{Flu})$ = fraction of flu-inflicted worlds in which you have a headache. How to calculate?

- Definition:

$$P(X|Y) = \frac{P(X, Y)}{P(Y)} = \frac{P(Y|X)P(X)}{P(Y)}$$

Corollary:

$$P(X, Y) = P(Y|X)P(X)$$

This is called **Bayes Rule**


Bayes' Rule

- $$P(\text{Headache}|\text{Flu}) = \frac{P(\text{Headache},\text{Flu})}{P(\text{Flu})}$$
$$= \frac{P(\text{Flu}|\text{Headache})P(\text{Headache})}{P(\text{Flu})}$$

Other cases:

- $$P(Y|X) = \frac{P(X|Y)P(Y)}{P(X|Y)P(Y)+P(X|\neg Y)P(\neg Y)}$$
- $$P(Y = y_i|X) = \frac{P(X|Y)P(Y)}{\sum_{i \in S} P(X|Y = y_i)P(Y=y_i)}$$
- $$P(Y|X, Z) = \frac{P(X|Y, Z)P(Y, Z)}{P(X, Z)} =$$
$$\frac{P(X|Y, Z)P(Y, Z)}{P(X|Y, Z)P(Y, Z)+P(X|\neg Y, Z)P(\neg Y, Z)}$$

Outline

- Probability Distributions
- Joint and Conditional Probability Distributions
- Bayes' Rule
- Mean and Variance 
- Properties of Gaussian Distribution
- Maximum Likelihood Estimation

Mean and Variance

- Expectation: The mean value, center of mass, first moment:

$$E_X[g(X)] = \int_{-\infty}^{\infty} g(x)p_X(x)dx = \mu$$

- N-th moment: $g(x) = x^n$
- N-th central moment: $g(x) = (x - \mu)^n$
- Mean: $E_X[X] = \int_{-\infty}^{\infty} xp_X(x)dx$
 - $E[\alpha X] = \alpha E[X]$
 - $E[\alpha + X] = \alpha + E[X]$
- Variance(Second central moment): $Var(x) = E_X[(X - E_X[X])^2] = E_X[X^2] - E_X[X]^2$
 - $Var(\alpha X) = \alpha^2 Var(X)$
 - $Var(\alpha + X) = Var(X)$

Expected value and average

Expectation determined by probability distribution. Arithmetic average determined by observed outcomes of trials.

Both are the same in our class because we assume we have sufficient data that models the distribution. So, the arithmetic average is a good estimate of the true expectation (Law of Large Numbers).

Variance and average:

Covariance:

Correlation:

Useful in EDA. If features are correlated, then data may have redundancy.

Uncorrelated vs Independent RV

Uncorrelated (Definition: $\text{Cov}(X, Y) = 0$.)

- Means no **linear relationship** between X and Y .
- Does **not** rule out non-linear dependence.
- Example: $X \sim U(-1, 1)$, $Y = x^2 \rightarrow$ Uncorrelated but dependent.

Independent (Definition: $P(X \in A, Y \in B) = P(X \in A)P(Y \in B)$)

- Stronger condition: knowing one variable gives **no information** about the other.
- Independence \Rightarrow Uncorrelated (if variances exist).

Key Differences

- Independence is a **stronger** property.
- Uncorrelated only removes **linear relationships**.
- Uncorrelated $\not\Rightarrow$ Independent.

ML Relevance

- Most ML models care about uncorrelatedness because they only model linear relationships.
- True independence is rarer and much harder to check, but it's what we assume in stronger probabilistic models

For Joint Distributions

- Expectation and Covariance:

- $E[X + Y] = E[X] + E[Y]$


- $cov(X, Y) = E[(X - E_X[X])(Y - E_Y(Y))] = E[XY] - E[X]E[Y]$

- $Var(X + Y) = Var(X) + 2cov(X, Y) + Var(Y)$

Recap

- Sum and Product Rule. Bayes Rule
- Expectation and arithmetic mean
- Variance
- Covariance
- Correlation
- Uncorrelated vs Independent r.v.


Outline

- Probability Distributions
- Joint and Conditional Probability Distributions
- Bayes' Rule
- Mean and Variance
- Properties of Gaussian Distribution 
- Maximum Likelihood Estimation

Gaussian Distribution

- Gaussian Distribution: $f(x|\mu, \sigma^2) = \frac{1}{\sqrt{2\pi\sigma^2}} e^{-\frac{(x-\mu)^2}{2\sigma^2}}$

Probability density function



Probability versus likelihood

Multivariate Gaussian Distribution

$$p(x|\mu, \Sigma) = \frac{1}{(2\pi)^{n/2} |\Sigma|^{1/2}} \exp\left\{-\frac{1}{2} (x - \mu)^\top \Sigma^{-1} (x - \mu)\right\}$$

- Moment Parameterization $\mu = E(X)$

$$\Sigma = \text{Cov}(X) = E[(X - \mu)(X - \mu)^\top]$$

- Mahalanobis Distance $\Delta^2 = (x - \mu)^\top \Sigma^{-1} (x - \mu)$
- Tons of applications (MoG, FA, PPCA, Kalman filter,...)

Properties of Gaussian Distribution

- The **linear transform** of a Gaussian r.v. is a Gaussian. Remember that no matter how x is distributed

$$E(AX + b) = AE(X) + b$$

$$\text{Cov}(AX + b) = ACov(X)A^T$$

this means that for Gaussian distributed quantities:

$$X \sim N(\mu, \Sigma) \rightarrow AX + b \sim N(A\mu + b, A\Sigma A^T)$$

- The **sum** of two independent Gaussian r.v. is a Gaussian

$$Y = X_1 + X_2, X_1 \perp X_2 \rightarrow \mu_y = \mu_1 + \mu_2, \Sigma_y = \Sigma_1 + \Sigma_2$$


- The **multiplication** of two Gaussian functions is another Gaussian function (although no longer normalized)

$$N(a, A)N(b, B) \propto N(c, C),$$

$$\text{where } C = (A^{-1} + B^{-1})^{-1}, c = CA^{-1}a + CB^{-1}b$$

Central Limit Theorem

Probability mass function of a **biased** dice



Let's say, I am going to get a sample from this pmf having a size of $n = 4$


$$S_1 = \{1,1,1,6\} \Rightarrow E(S_1) = 2.25$$

$$S_2 = \{1,1,3,6\} \Rightarrow E(S_2) = 2.75$$

⋮

$$S_m = \{1,4,6,6\} \Rightarrow E(S_m) = 4.25$$

- According to CLT, if you sample enough from any distribution with finite variance you will get an approximate Gaussian distribution.
- No matter what the population looks like, the average of many samples looks Normal.
- Explains **why the Normal distribution is everywhere** in statistics and ML.



Prob vs Likelihood

*Probability predicts data from parameters.
Likelihood evaluates parameters from data.*

•Probability

- Forward direction: given parameters, what's the chance of data?
- $P(data | \theta)$
- Example: *If the coin has bias $\theta = 0.7$, what's the probability of 8 heads in 10 tosses?*
- Varies with **different outcomes** of data.

•Likelihood

- Reverse view: given data, how plausible are parameters?
- $L(\theta | data) \propto P(data | \theta)$
- Example: *I observed 8 heads in 10 tosses. How likely is it that the coin's bias is $\theta = 0.7$?*
- Varies with **different parameter values**.

Prob vs Likelihood



Prob vs Likelihood



Outline

- Probability Distributions
- Joint and Conditional Probability Distributions
- Bayes' Rule
- Mean and Variance
- Properties of Gaussian Distribution
- Maximum Likelihood Estimation ←

Maximum Likelihood Estimation

- Probability: inferring probabilistic quantities for data given fixed models (e.g. prob. of events, marginals, conditionals, etc).
- Statistics: inferring a model given fixed data observations (e.g. clustering, classification, regression).

Main assumption:

Independent and identically distributed random variables
i.i.d

Maximum Likelihood Estimation

For Bernoulli (i.e. flip a coin):

Objective function: $P(x^{i}|\theta) = \theta^{x^{i}}(1-\theta)^{1-x^{i}}$ $x^{i} \in \{0,1\}$ or {head, tail}

$$L(\theta|X) = L(\theta|X = x^{1}, X = x^{2}, X = x^{3}, \dots, X = x^{n})$$

i.i.d assumption

$$L(\theta|X) = \prod_{i=1}^n P(x^{i}|\theta)$$

$$L(\theta|X) = \prod_{i=1}^n P(x^{i}|\theta) = \prod_{i=1}^n \theta^{x^{i}}(1-\theta)^{1-x^{i}}$$

$$\begin{aligned} L(\theta|X) &= \theta^{x^{1}}(1-\theta)^{1-x^{1}} \times \theta^{x^{2}}(1-\theta)^{1-x^{2}} \dots \times \theta^{x^{n}}(1-\theta)^{1-x^{n}} = \\ &= \theta^{\sum x^{i}}(1-\theta)^{\sum(1-x^{i})} \end{aligned}$$

We don't like multiplication, let's convert it into summation

What's the trick?

Take the log

$$L(\theta|X) = \theta^{\sum x^{i}} (1 - \theta)^{\sum (1-x^{i})}$$

$$\log L(\theta|X) = l(\theta|X) = \log(\theta) \sum_{i=1}^n x^{i} + \log(1 - \theta) \sum_{i=1}^n (1 - x^{i})$$

How to optimize θ ?

$$\frac{\partial l(\theta|X)}{\partial \theta} = 0 \quad \frac{\sum_{i=1}^n x^{i}}{\theta} - \frac{\sum_{i=1}^n (1 - x^{i})}{1 - \theta} = 0$$

$$\theta = \frac{1}{n} \sum_{i=1}^n x^{i}$$